

CORRELATION OF A 3,200 YEAR OLD TEPHRA IN ICE CORES FROM VOSTOK AND SOUTH POLE STATIONS, ANTARCTICA

J.M. Palais¹, P.R. Kyle², E. Mosley-Thompson³ and E. Thomas⁴

Abstract. Tephra layers in two Antarctic ice cores are correlated on the basis of their chemical compositions and estimated ages. We believe the tephra was produced about 3,200 years ago from a major explosive eruption of Candlemas Island in the South Sandwich Islands. This is the first time that tephra layers have been correlated between two widely separated ice cores and demonstrates that tephra layers may serve as stratigraphic markers for correlating, and perhaps in some cases, dating Antarctic ice cores.

Introduction

Studies of ice core samples from Greenland and Antarctica provide detailed information about past climate and the chemical composition of the atmosphere, for at least the last 150,000 years (Dansgaard et al., 1981, 1982; Lorius et al., 1979, 1985; Oeschger, 1985). The preservation of volcanic deposits (both soluble acids and insoluble volcanic ash particles) from both local and distant volcanic sources in Antarctic and Greenland ice (Hammer et al., 1981; Kyle et al., 1982, 1984; Hammer, 1984; Palais, 1985; De Angelis et al., 1985; Delmas et al., 1985) is of particular interest.

Eruptions from Antarctic volcanoes in the McMurdo Volcanic Group and Marie Byrd Land (Keys et al., 1977; Kyle et al., 1981) and volcanoes in the South Sandwich Islands (Nishio et al., 1984 a,b) are recorded principally as visible tephra (volcanic ash) layers in the ice cores, whereas eruptions originating from volcanoes in the more distant regions of Indonesia, South America, and New Zealand have produced only layers of elevated acidity (e.g. H₂SO₄), recognized by detailed analyses of the chemical composition of the ice (Delmas et al., 1985).

In Greenland, eruptions of nearby Icelandic volcanoes have produced well-defined layers of elevated acidity, identified by increases in the D.C. electrical conductivity of the ice, but no insoluble volcanic ash particles have ever been identified (Hammer et al., 1981; Hammer, 1984). The only well confirmed evidence to date, for long-distance (> several thousand kilometers) transport of volcanic ash particles to the polar ice sheets, in either hemisphere, is the study by De Angelis et al. (1985), in which volcanic ash particles originating in the 1982 eruption of El Chichon, Mexico, were found in surface snow of the Greenland ice sheet. Further evidence that tephra can be transported far from source, and is therefore useful for correlation purposes, is the study by Kyle and Seward (1984), in which tephra from New Zealand was identified in deep-sea piston cores from the South Pacific sector of the Southern Ocean near the Balleny Islands.

In this paper we report the first correlation of tephra layers between two ice cores in Antarctica. The visible tephra layers in ice cores from Vostok Station (78° 28' S, 106° 48' E) and South Pole Station (90° S), have been correlated on the basis

of their chemical composition and estimated age. We believe that the tephra originated in a major explosive eruption from Candlemas Island in the South Sandwich Islands about 3,200 years ago.

Sampling and Analysis

Ice samples containing the two tephra layers were melted and filtered onto Nucleopore™ (0.4 μm diameter pore size) filters in a Class 100 clean room. A section from each filter was examined by scanning electron microscope (SEM) and energy-dispersive x-ray analysis (EDAX). A polished microprobe section of each sample was prepared by washing the particles off of the filter and mounting them in epoxy.

The energy-dispersive x-ray analyses use "ZAF" corrections to calculate oxide percents (normalized to 100%), from element peak intensities and theoretical standards. The accuracy of the analyses was checked against glass and mineral standards and the precision of the analyses was determined by repeated analyses of these same standards. The microprobe analyses were corrected using the Bence-Albee matrix correction procedure (Bence and Albee, 1968) and glass standards were used to check analytical precision and accuracy.

The microprobe analyses are more precise than the EDAX analyses, as can be seen by comparing the standard deviations of the analyses (Table 1).

Results

Vostok Tephra

The Vostok tephra is the informal name proposed by Kyle et al. (1982) for a 0.05 m thick, diffuse tephra layer found at 100.8 m in an ice core drilled in 1979 at Vostok Station in East Antarctica (Figure 1). The depth/age relationship (Parker et al., 1982) for the Vostok core suggests an age of about 3,200 a B.P. The Vostok tephra is composed of 55 % lithic material (mainly well-rounded, brown, cryptocrystalline glass), 40 % clear glass shards and 5 % crystals (mainly feldspar) (Figure 2 c,d).

Kyle et al. (1984) considered three possible source regions (the South Shetland Islands, the South Sandwich Islands and the Southern Andes) for the Vostok tephra and concluded that the South Sandwich Islands were the most likely source. Rocks from the South Shetland Islands and the Southern Andes have higher total alkali content and lower iron contents, respectively, than the Vostok tephra. The Vostok tephra (Table 1, sample 3,4) is most similar in composition to an aphyric andesite from the northern part of Candlemas Island in the South Sandwich Islands, which is one of the youngest of the South Sandwich Islands (Table 1, sample 5) (Baker, 1978; Tomblin, 1979). Lavas from Candlemas Island are exceptionally low in potash (K₂O), a characteristic feature of the Vostok tephra.

Grain size analyses of the Vostok tephra indicate initial deposition of coarse particles (mean diameter of about 45 μm) followed by deposition of much finer particles (mean of about 7 μm), composing the bulk of the tephra (Figure 3). The mass concentration of tephra in melt water samples gradually increases from a background of about 1 μg/ml to a peak of about 14 μg/ml. The original thickness of the tephra is unknown as the diffuse nature of the Vostok tephra in the core makes it difficult to calculate a mass accumulation rate (mg/cm²/1000a), as is done in deep sea cores.

¹ Graduate School of Oceanography, Univ. of Rhode Island, Narragansett, R.I.

² Dept. of Geoscience, New Mexico Institute of Mining and Technology, Socorro, N.M.

³ Byrd Polar Research Center, Ohio State University, Columbus, Ohio

⁴ Lamont-Doherty Geological Observatory, Palisades, N.Y.

Copyright 1987 by the American Geophysical Union.

Paper number 7L7157.

0094-8276/87/007L-7157\$03.00




Fig. 1. Map of Antarctica showing core locations and probable paths of atmospheric transport from the South Sandwich Islands to the Antarctic ice sheet.

South Pole Tephra

In 1983/84 a 361 m ice core was drilled at the South Pole with a distinct tephra layer (3mm thick) at 303.44 m. The tephra is composed primarily of vesicular glass shards and lithic material with a small percentage of crystals, including quartz, pyroxene and plagioclase of intermediate composition (Figure 2 a,b). Major element analyses of the glass shards in the South Pole tephra (Table 1, analyses 1,2) compare well

Table 1. Major element analyses (wt. %) of tephra from the South Pole and Vostok cores, and a rock from the South Sandwich Islands.

	1	2	3	4	5
n	5	11	9	37	
SiO ₂	59.98(1.21)†	60.1(3.02)	60.56(0.42)	59.73(1.79)	60.9
TiO ₂	0.78(0.9)	0.3(0.22)	0.74(0.07)	0.90(0.16)	0.95
Al ₂ O ₃	16.03(0.94)	17.6(3.40)	14.92(0.82)	14.84(0.66)	14.8
FeO*	7.37(1.14)	9.0(3.79)	9.15(0.57)	10.25(1.23)	8.34
MnO	0.16(0.8)	0	0.23(0.04)	0.15(0.12)	0.11
MgO	1.71(0.28)	3.5(1.91)	2.39(0.34)	2.44(0.48)	2.35
CaO	6.79(0.42)	5.5(1.35)	6.72(0.26)	6.64(0.76)	6.09
Na ₂ O	5.94(0.83)	3.6(0.89)	3.31(0.38)	4.46(0.57)	3.67
K ₂ O	0.39(0.09)	0.1(0.15)	0.44(0.05)	0.63(0.19)	0.39
P ₂ O ₅	0.08(0.09)	0.3(0.13)	n.d.	n.d.	0.12
Total	99.23	100	98.46	100.04	97.72

Key:

* FeO, total Fe as FeO;

† one standard deviation n.d. indicates not detected; n is number of samples.

1. South Pole Tephra- Electron microprobe analyses, Univ. of Rhode Island, (Analyst, J. Palais).
2. South Pole Tephra-Energy-dispersive x-ray analysis, Ohio State Univ. (Analyst, T. Leonardi).
3. Vostok Tephra- Electron microprobe analysis, Los Alamos National Laboratory (Analyst, J. Palais) (Kyle et al., 1984).
4. Vostok Tephra- Energy dispersive x-ray analysis, Arizona State University (Analyst, E. Thomas) (Kyle et al., 1984).
5. Candlemas Island-Aphyric andesite, Cauldron Lake lava, northern Candlemas Island. Wet chemical analysis (Tomblin, 1979).




Fig. 2. Scanning electron micrographs of tephra from the South Pole and Vostok ice cores. a) South Pole tephra blocky, moderately vesicular, partially altered sherd with shallow spherical vesicles. Fine adhering dust. 2400 x; 1 cm=4 μm. b) South Pole tephra highly vesicular, fresh sherd with delicate square to rectangular vesicles. 2400x; 1 cm=4 μm. c) Vostok tephra subrounded, blocky, moderately vesicular sherd, partially altered, with shallow spherical vesicles. Fine adhering dust. Compare with a). 1110 x; 1 cm= 9 μm d) Vostok tephra blocky shards with fine adhering dust, few vesicles. 1110 x; 1cm= 9 μm.




Fig. 3. Depth variation of mass concentration of tephra in a 0.10 m section of the Vostok core. Histograms show the grain size distribution of tephra in the samples that were analyzed for mass concentration. Sulfate and nitrate data from Kyle et al. (1982).

with those of the Vostok tephra (Table 1, analyses 3,4), which suggests that they might be from the same eruption. This is further supported by the estimated age of the ice enclosing these layers, as discussed below.

Depth-Age Relationships

There is no absolute dating of either the South Pole or Vostok cores, but two lines of evidence suggest that the tephra layers (Vostok tephra-100m; South Pole tephra-300m) are coeval. First, a rough estimate of the age of these two layers is made by assuming a similar depth-density profile for the two locations and accumulation rates which differ by a factor of about three (Vostok 2.0-2.5 g/cm²/a (Young et al., 1982; Lorius et al., 1985); South Pole 7-9 g/cm²/a (Jouzel et al., 1983)). Secondly, our estimated depth-age profile for the South Pole core (Figure 4), based on depth-age estimates of Schwander (1984), Mosley-Thompson (1980) and Kuivinen et




Fig. 4. Estimated depth-age relationship for the South Pole core based on data from Schwander (1984), Mosley-Thompson (1980) and Kuivinen et al (1982).

al. (1982), suggests an age of 3,250 a B.P. at 300 m in the South Pole core. This compares well with the estimated age (3,200 a B.P.) for the Vostok tephra (Parker et al., 1982).

Discussion

Tephra similar in composition to that in the Vostok and South Pole cores also occurs in layers at the surface in blue ice areas (the Meteorite Ice Field) of the Yamato Mountains, Queen Maud Land, East Antarctica (Figure 1) (Nishio et al., 1984; Katsushima et al., 1984). Although the ages of the tephra layers in the Yamato Mountains are not well constrained, they are of the correct order of magnitude to be correlative with the Vostok and South Pole tephra. ³⁶Cl analyses of ice associated with these tephra layers, suggest that the age of the ice in the Meteorite Ice Field of the Yamato Mountains is less than one ³⁶Cl half-life (3×10^5 a) (Nishiizumi et al., 1979). The ¹⁴C terrestrial age of chondrite Yamato 75102, collected near tephra layer C-32, indicates an age of $3.0 \pm 1.2 \times 10^3$ a (Nishio, personal communication, 1985). The terrestrial ages of four other Yamato meteorites are believed to be in the range of 3,000-22,000 a. Calculation of flow lines in the Meteorite Ice Fields gives an age for the ice upstream of Motoi Nunatak, near tephra layer C-32, in the range from one to several thousand years old (Nishio et al., 1984).

Tephra layers in piston cores from the South Atlantic Ocean (Ninkovich et al., 1964; Federman et al., 1982) were believed to be erupted from the South Sandwich Islands within the last 30,000 years. Based on a sedimentation rate of 8 cm/1000 a, a tephra layer at 0.20 m in core V 14-56 would be 2,500 years old (Federman et al., 1982). This is probably a minimum estimate, however, since erosion by Antarctic Bottom Water may have removed some of the sediments in the core (Ledbetter and Ciesielski, 1986). The tephra has a similar composition and age to the tephra layers from Vostok, South Pole and from the Yamato Mountains and may correlate with them.

Conclusions

These analyses and inferences are consistent with the conclusion that the tephra layers in the South Pole and Vostok cores are coeval and represent deposition from a major explosive volcanic eruption in the South Sandwich Islands (probably from Candlemas Island) about 3,200a B.P. For the first time tephra layers from two widely separated ice cores have been correlated, which reinforces the utility of tephrochronology for ice core studies. This tephra layer provides an excellent stratigraphic marker for Antarctic ice cores and should be sought at other locations. Furthermore, a detailed study of the dispersal of this tephra layer may provide information on the transport of atmospheric particles in this part of the southern hemisphere. The occurrence of an apparently widely dispersed tephra layer in East Antarctica, possibly originating from the South Sandwich Islands, is consistent with the paths of atmospheric transport proposed by Alt et al. (1959), Carleton (1981) and Thompson and Mosley-Thompson (1982).

Acknowledgements. We would like to thank the Polar Ice Coring Office (PICO), University of Nebraska for drilling the Vostok and South Pole ice cores. We also wish to thank Grant Heiken for use of the Electron Microprobe at the Los Alamos National Laboratory and Peter Busek for use of the Scanning Electron Microscope at the Arizona State University. Dave Mann (Mann Petrographic Inc.) made the microprobe mounts of both tephra samples. Tony Leonardi of the Ohio State University and David Browning of the University of Rhode Island assisted with the SEM and EDAX analyses of the South Pole tephra. This work was supported by the National Science Foundation grant DPP-8021402, DPP-8519122

References

- Alt, J., P. Astapenko, N.J. Ropar, Jr., Some aspects of the Antarctic atmospheric circulation in 1958, *IGY General Report Series 4*, 1959.
- Baker, P.E. The South Sandwich Islands: III Petrology of the volcanic rocks, Brit. Antarct. Surv. Sci. Rpt. 93, 1-34, 1978.
- Bence, A.E. and A.L. Albee, Empirical correction factors for the electron microanalysis of silicates and oxides, *J. Geol.*, **76**, 382-403, 1968.
- Carleton, A.M., Monthly variability of satellite-derived cyclone activity for the southern hemisphere winter, *J. Climatol.*, **1**, 21-38, 1981.
- Dansgaard, W., Ice core studies: dating the past to find the future, *Nature*, **290**, 360-361, 1981.
- Dansgaard, W., H.O. Clausen, N. Gundestrup, C.U. Hammer, S.J. Johnsen, P.M. Kristinsdottir, and N. Reeh, A new Greenland deep ice core, *Nature*, **218**, 1273-1277, 1982.
- De Angelis M., L. Fehrenbach, C. Jéhanno, and M. Maurette, Micrometre-sized volcanic glasses in polar ices and snows, *Nature*, **317**, 52-54, 1985.
- Delmas, R.J., M. Legrand, A.J. Arístarain and F. Zanolini, Volcanic deposits in Antarctic snow and ice, *J. Geophys. Res.*, **90**, 12,901-12,920, 1985.
- Federman, A.W., N.D. Watkins and H. Sigurdsson, Scotia Arc volcanism recorded in abyssal piston cores downwind from the islands, in *Antarctic Geoscience*, edited by C. Craddock, pp. 223-228, University of Wisconsin Press, Madison, Wisc., 1982.
- Hammer, C.U., H.B. Clausen, W. and Dansgaard, Greenland ice sheet evidence of post-glacial volcanism and its climatic impact, *Nature*, **288**, 230-235, 1981.
- Hammer, C.U., Traces of Icelandic eruptions in the Greenland ice sheet, *Jökull*, **34**, 51-65, 1984.
- Jouzel, J., L. Merlivat, J.R. Petit and C. Lorius, Climatic information over the last century deduced from a detailed isotopic record in the South Pole snow, *J. Geophys. Res.*, **88**, 2693-2703, 1983.
- Katsushima, T., F. Nishio, H. Ohmae, M. Ishikawa and S. Takahashi, Composition of dirt layers in the bare ice areas near the Yamato Mountains in Queen Maud Land and the Allan Hills in Victoria Land, Antarctica, *Mem. Nat. Inst. Polar Res. Spec. Issue 34*, 174-187, 1984.
- Kuivinen, K.C., B.R. Koci, G.W. Holdsworth and A.J. Gow, South Pole ice core drilling, 1981-1982, *Antarct. J. U.S.*, **17**, 89-91, 1982.
- Kyle, P.R., P.A. Jezek, E. Mosley-Thompson and L.G. Thompson, Tephra layers in the Byrd Station ice core and the Dome C ice core, Antarctica and their climatic importance, *J. Volcanol. Geotherm. Res.*, **11**, 29-39, 1981.
- Kyle, P.R., J. M. Palais and R. Delmas, The volcanic record of Antarctic ice cores: Preliminary results and potential for future investigations, *Annals Glaciol.*, **3**, 172-179, 1982.
- Kyle, P.R., J. Palais, J. and E. Thomas, The Vostok tephra—An important englacial stratigraphic marker?, *Antarct. J. U.S.*, **19**, 64-65, 1984.
- Kyle, P.R. and D. Seward, Dispersed rhyolitic tephra from New Zealand in deep-sea sediments of the Southern Ocean, *Geology*, **12**, 487-490, 1984.
- Ledbetter, M.T. and P. F. Ciesielski, Post-Miocene disconformities and paleoceanography in the Atlantic sector of the Southern Ocean, *Paleogeograph., Paleoclimatol., Paleoecol.*, **52**, 185-214, 1986.
- Lorius, C., L. Merlivat, J. Jouzel, and M. Pourchet, A 30,000-year isotope climatic record from Antarctic ice, *Nature*, **280**, 644-648, 1979.
- Lorius, C., J. Jouzel, C. Ritz, L. Merlivat, N.I. Barkov, Y.S. Korotkevich and V.M. Kotlyakov, A 150,000-year climatic record from Antarctic ice, *Nature*, **316**, 591-596, 1985.
- Mosley-Thompson, E. 911 Years of Microparticle Deposition at the South Pole: A Climatic Interpretation, Ohio State Univ. Inst. Polar Studies, Ohio State Univ., Rpt. 73, 1980.
- Ninkovich, D., B. C. Heezen, J.R. Conolly and L.H. Burkle, South Sandwich tephra in deep-sea sediments, *Deep Sea Res.*, **11**, 605-619, 1964.
- Nishiizumi, K., L.R. Arnold, D. Elmore, R.D. Ferraro, H.E. Gove, R.C. Finkel, R.P. Beukens, K.H. Chang and L.R. Kilius, ^{36}Cl and ^{53}Mn in Antarctic meteorites and ^{10}Be - ^{36}Cl dating of Antarctic ice, *Earth Planet. Sci. Lett.*, **45**, 285-292, 1979.
- Nishio, F., T. Katsushima and H. Ohmae, Volcanic ash layers in bare ice areas near the Yamato Mountains, Dronning Maud Land and the Allan Hills, Victoria Land, Antarctica., *Annals Glaciol.*, **7**, 34-41, 1984a.
- Nishio, F., T. Katsushima, H. Ohmae, M. Ishikawa and S. Takahashi, Dirt layers and atmospheric transportation of volcanic glass in bare ice areas near the Yamato Mountains in Queen Maud Land and the Allan Hills in Victoria Land in Antarctica, *Nat. Inst. Polar Res. Spec. Issue*, **34**, 160-173, 1984b.
- Oeschger, H. The contribution of ice core studies to the understanding of environmental processes, in *Greenland Ice Core: Geophysics, Geochemistry, and the Environment*, edited by C.C. Langway, Jr., H. Oeschger, and W. Dansgaard, 9-17, Am. Geophys. Union, Geophysical Monograph 33, Wash. D.C., 1985.
- Palais, J.M., Tephra layers and ice chemistry in the Byrd-Station ice core, Antarctica, PhD. dissertation, 516 pp. The Ohio State Univ., Columbus, Aug. 1985.
- Parker, B.C., E.J. Zeller and A.J. Gow, Nitrate fluctuation in Antarctic snow and firn: Potential sources and mechanisms of formation, *Annals Glaciol.*, **3**, 243-248, 1982.
- Schwander, J. Luftabschluss im eis von Gronland und der Antarktis, messung der elektrischen leit fähigkeit von eisproben für klimatologische anwendungen, PhD. dissertation, Univ. of Bern, Switz., 1984.
- Thompson, L.G. and E. Mosley-Thompson, Spatial distribution of microparticles within Antarctic snow-fall, *Annals Glaciol.*, **3**, 300-306, 1982.
- Tomblin, J.F., The South Sandwich Islands: II The Geology of Candlemas Island, Brit. Antarct. Surv. Sci. Rpt. 92, 1-33, 1979.
- Watkins, N.D. and T.C. Huang, Tephra in abyssal sediments east of North Island, New Zealand, chronology, paleowind velocity and paleoexplosivity, *N.Z. J. Geol. Geophys.*, **20**, 179-198, 1977.
- Young, N.W., M. Pourchet, V.M. Kotlyakov, P.A. Korolev and M.B. Dyugorov, Accumulation distribution in the IAGP area, Antarctica: 90°E-150°E, *Annals Glaciol.*, **3**, 333-338, 1982.

J. Palais, Graduate School of Oceanography, U.R. I., Narragansett, R.I., 02882.

P. Kyle, Department of Geoscience, New Mexico Institute of Mining and Technology, Socorro, N.M., 87801.

E. Mosley-Thompson, Byrd Polar Research Center, Ohio State University, Columbus, Ohio, 43210.

E. Thomas, Lamont-Doherty Geological Observatory, Palisades, N.Y. 10964.

(Received March 17, 1987;
accepted April 27, 1987.)